



**University of
Zurich**^{UZH}

**Zurich Open Repository and
Archive**

University of Zurich
University Library
Strickhofstrasse 39
CH-8057 Zurich
www.zora.uzh.ch

Year: 2011

A large-scale forest fragmentation experiment: the stability of altered forest ecosystems project

Ewers, R M ; Didham, R K ; Fahrig, L ; Ferraz, G ; Hector, A ; Holt, R D ; Kapos, V ; Reynolds, G ; Sinun, W ; Snaddon, J L ; Turner, E C

Abstract: Opportunities to conduct large-scale field experiments are rare, but provide a unique opportunity to reveal the complex processes that operate within natural ecosystems. Here, we review the design of existing, large-scale forest fragmentation experiments. Based on this review, we develop a design for the Stability of Altered Forest Ecosystems (SAFE) Project, a new forest fragmentation experiment to be located in the lowland tropical forests of Borneo (Sabah, Malaysia). The SAFE Project represents an advance on existing experiments in that it: (i) allows discrimination of the effects of landscape-level forest cover from patch-level processes; (ii) is designed to facilitate the unification of a wide range of data types on ecological patterns and processes that operate over a wide range of spatial scales; (iii) has greater replication than existing experiments; (iv) incorporates an experimental manipulation of riparian corridors; and (v) embeds the experimentally fragmented landscape within a wider gradient of land-use intensity than do existing projects. The SAFE Project represents an opportunity for ecologists across disciplines to participate in a large initiative designed to generate a broad understanding of the ecological impacts of tropical forest modification.

DOI: <https://doi.org/10.1098/rstb.2011.0049>

Posted at the Zurich Open Repository and Archive, University of Zurich

ZORA URL: <https://doi.org/10.5167/uzh-52549>

Journal Article

Accepted Version

Originally published at:

Ewers, R M; Didham, R K; Fahrig, L; Ferraz, G; Hector, A; Holt, R D; Kapos, V; Reynolds, G; Sinun, W; Snaddon, J L; Turner, E C (2011). A large-scale forest fragmentation experiment: the stability of altered forest ecosystems project. *Philosophical Transactions of the Royal Society B: Biological Sciences*, 366(1582):3292-3302.

DOI: <https://doi.org/10.1098/rstb.2011.0049>

**A large-scale forest fragmentation experiment: the Stability of Altered
Forest Ecosystems Project**

**Robert M. EWERS^{1*}, Raphael K. DIDHAM^{2,3,4}, Lenore FAHRIG⁵, Gonçalo
FERRAZ^{6,7}, Andy HECTOR⁸, Robert D. HOLT⁹, Valerie KAPOS^{10,11}, Glen
REYNOLDS¹², Waidi SINUN¹³, Jake L. SNADDON^{8,11} and Edgar C. TURNER^{1,11}**

¹ Imperial College London, Silwood Park Campus, Buckhurst Road, Ascot, Berkshire SL5 7PY,
UK

² School of Animal Biology, The University of Western Australia, 35 Stirling Highway, Crawley
WA 6009, Australia

³ CSIRO Ecosystem Services, Centre for Environment and Life Sciences, Underwood Avenue,
Floreat WA 6014, Australia

⁴ School of Biological Sciences, University of Canterbury, Private Bag 4800, Christchurch, New
Zealand

⁵ Ottawa-Carleton Institute of Biology, Carleton University, Ottawa, Ontario, Canada K1S 5B6

⁶ Biological Dynamics of Forest Fragments Project, Smithsonian Tropical Research
Institute/Instituto Nacional de Pesquisas da Amazônia, 69011 Manaus AM, Brazil

⁷ Smithsonian Tropical Research Institute, Apartado 0843-03092, Balboa, Republic of Panama

⁸ Institute of Evolutionary Biology and Environmental Studies, University of Zurich,
Winterthurerstrasse 190, CH-8057 Zurich, Switzerland

⁹ Department of Biology, University of Florida, 111 Bartram, P.O. Box 118525, Gainesville, FL
32611, USA

¹⁰ United Nations Environmental Programme (UNEP) World Conservation Monitoring Centre
(WCMW), 219 Huntingdon Road, Cambridge CB3 0DL, UK

¹¹ Department of Zoology, University of Cambridge, Downing Street, Cambridge CB2 3EJ, UK

¹² The Royal Society South East Asian Rainforest Research Programme (SEARRP), Danum Valley Field Centre, PO Box 60282, Lahad Datu 91112, Sabah, Malaysia

¹³ Conservation and Environmental Management Division, Yayasan Sabah Group, Menara Tun Mustapha, Kota Kinabalu, Sabah 88817, Malaysia

* Corresponding Author. Email: r.ewers@imperial.ac.uk;

Running header: The SAFE Project

Abstract

Opportunities to conduct large-scale field experiments are rare, but provide a unique prospect to reveal the complex processes that operate within natural ecosystems. Here, we review the design of existing, large-scale forest fragmentation experiments. Based on this review, we develop a design for the Stability of Altered Forest Ecosystems (SAFE) Project, a new forest fragmentation experiment to be located in the lowland tropical forests of Borneo (Sabah, Malaysia). SAFE represents an advance on existing experiments in that it: (1) allows discrimination of the effects of landscape-level forest cover from patch-level processes; (2) is designed to facilitate the unification of a wide range of data types collected on ecological patterns and processes that operate over a wide range of spatial scales; (3) has greater replication than existing experiments; (4) incorporates an experimental manipulation of riparian corridors; and (5) embeds the experimentally fragmented landscape within a wider gradient of land use intensity than existing projects. The SAFE Project represents an opportunity for ecologists across

disciplines to participate in a large initiative designed to generate a broad understanding of the ecological impacts of tropical forest modification.

Keywords: Biological Dynamics of Forest Fragments Project; Calling Lake Fragmentation Experiment; deforestation; hierarchical sampling design; Savannah River Site Corridor Experiment; Wog Wog Habitat Fragmentation Experiment

1. INTRODUCTION

Habitat fragmentation is one of the central issues in conservation biology [1] and has been a source of considerable scientific debate since the early application of island biogeography theory [2] to terrestrial habitat islands and the design of nature reserves [3-4]. These debates led directly to the establishment of the Minimum Critical Size of Ecosystems experiment in the Brazilian Amazon [5-6], now known as the Biological Dynamics of Forest Fragments Project (BDFFP). This visionary experiment has probably had the single greatest impact on the general understanding of the ecological impact of forest fragmentation and is routinely cited in the conservation literature around the world [7-8].

The BDFFP heralded a new approach to the study of habitat fragmentation, lifting it from one based almost solely on observational studies, to one based on experimentation. The implications for scientific advancement were tremendous: the use of statistically based before-after-control-impact sampling provides much stronger inference than observational studies relying on a space-for-time substitution to act as a control [9].

Since the establishment of the BDFFP, there have been at least 20 more experimental tests of habitat fragmentation [10], based mostly in grasslands or fields, but just one of these has matched the BDFFP in terms of the size and number of fragments [11].

Large-scale experiments may be the only way to determine the responses of forest systems to global change [12]. They generate results that are comparable to those arising from observational studies [13] which ensures that large-scale experiments have real-world practical relevance. However, the opportunities for large-scale, replicated and controlled landscape experiments are rare. When they are undertaken, they are invariably time-consuming and costly to establish, and consequently there will always be a premium placed on maximising the long-term scientific pay-off from this initial investment. Consequently, any such experiments should address a wide range of ecological questions [10]. Moreover, just as the BDFFP was designed to address a central, policy-relevant issue about reserve design, it is incumbent upon researchers establishing new fragmentation projects to ensure that they address not only policy questions that are relevant at the time of project establishment, but also for the foreseeable future.

Focusing on four large-scale forest fragmentation experiments located on three different continents, we summarise the key features of existing experimental designs and assess their ability to address emerging issues in landscape ecology. Based on this review, we develop and scrutinise the design for the Stability of Altered Forest Ecosystems (SAFE) Project, a new rainforest fragmentation experiment being established in the wet tropical forests of Malaysian Borneo.

96

97 **2. AN OVERVIEW OF EXPERIMENTAL DESIGNS FOR FOREST**

98 **FRAGMENTATION STUDIES**

99 Habitat area is probably the single most influential variable that can be manipulated in a
100 fragmentation experiment, and has formed the backbone of the designs at the BDFFP, the
101 Calling Lake Fragmentation Experiment, and the Wog Wog Habitat Fragmentation
102 Experiments. All three experiments used replicates of a small number of fragment size
103 categories, rather than opting for a more extensive, continuous gradient of unreplicated
104 fragment sizes. In all cases, fragment sizes were distributed on a log-scale. The BDFFP
105 initially planned four size categories on a \log_{10} scale (10^0 , 10^1 , 10^2 and 10^3 ha), but the
106 largest of these was never isolated. Calling Lake followed a similar design, but added an
107 intermediate fragment size of 40 ha. Wog Wog, however, had a much smaller forest area
108 available for the experiment and used fragment sizes of 0.25, 0.875 and 3.062 ha.

109

110 Surprisingly, given the island biogeography framework on which these experiments were
111 initially designed, isolation (or conversely connectivity) has seldom been built into large
112 scale fragmentation experiments. Only the Savannah River Site Corridor Experiment has
113 manipulated isolation distances, with fragments separated from each other by distances of
114 64 to 384 m [14]. Two other experiments used a coarser approach to manipulate isolation
115 by introducing a ‘connected’ versus ‘isolated’ treatment [11, 15]. A second experiment
116 at the Savannah River site was focused around issues of fragment connectivity, with
117 connected *vs* unconnected patches forming the central basis of the experimental design
118 [15]. Calling Lake also addressed connectivity in this manner, with connected fragments

119 cleared on just three of the four sides leaving the fragment directly connected to
120 continuous forest [11]. The unconnected fragments at Calling Lake are separated by a
121 fixed distance of 200 m from continuous forest, which is similar to the average (but
122 variable) distance of isolation at the BDFFP. By contrast, the fragments at Wog Wog are
123 as close as 50 m to each other and approximately the same distance from continuous
124 forest [16]. Forest regeneration in the matrix habitat at the BDFFP has allowed
125 inadvertent tests of connectivity, in that matrix regrowth has connected fragments to the
126 surrounding continuous forest over time [17].

127
128 The potential for ecological patterns to be obscured by environmental variability means
129 that the replication and spatial interspersions of fragments that differ in area should be
130 maximised [16]. All four forest fragmentation projects used a blocked experimental
131 design and a similar number of replicate fragments per block to allow for this. However,
132 the scale of landscape manipulations is often constrained by land availability, as well as
133 fiscal and logistical considerations, setting a limit on the amount of replication that can be
134 practicably achieved. Savanna River employed five or six replicate blocks (a total of 27
135 and 30 fragments) for the two experiments respectively, Wog Wog employed four
136 replicate blocks of fragments (a total of 12 fragments), whereas the BDFFP (11
137 fragments) and Calling Lake (12 isolated and 8 connected fragments) both used three
138 blocks. However, the Calling Lake landscape required that one of the blocks have a 100-
139 ha fragment that is widely separated from the others, and blocks in the BDFFP likewise
140 vary in terms of the number and size range of fragments: Dimona (2×1-ha, 1×10-ha and

1×100-ha fragments), Porto Alegre (1×1-ha, 1×10-ha and 1×100 ha fragments) and Esteio (2×1-ha and 2×10-ha fragments).

2.1 Emerging questions in landscape ecology and the ability of existing experiments to address them

Most early habitat fragmentation studies focussed on patch-level patterns and processes such as area, isolation and edge effects. These analyses typically ignored the wider landscape context within which the fragments themselves were embedded. The ongoing coupling of fragments to a surrounding landscape, which contains its own distinct community and suite of ecosystem processes, provides a generic source of differences from the expectations of fragments as true ‘islands’, surrounded by a matrix that is solely viewed as a barrier to dispersal. One problem that has thus emerged recently in analyses of fragmentation studies is that the patterns these early studies were ascribing to patch variables may have been confounded due to strong correlations between the patch variables and total amount of habitat area in the landscape [18-19]. Unless duly considered, observed fragmentation impacts could then, in some cases, be incorrectly attributed to patch rather than landscape variables. There can be no doubt that reducing the amount of forest in a landscape will have direct impacts on biodiversity, but very few studies have carefully teased apart the effects of habitat amount from the effects of the spatial configuration of that habitat [19]. Rigorously testing the relative impacts of area and configuration is challenging and requires identification of paired landscapes of the same size and with the same total amount of habitat, but with distinctively different habitat configurations.

164

165 A related issue revolves around the relative importance of forest cover patterns that are
166 measured at the scale of a single fragment, vs. those that are measured at the scale of a
167 landscape. Donovan et al. [20] provided one of the first simultaneous analyses of patch
168 and landscape features and found that edge effects (a patch-level pattern) varied
169 according to the amount of forest in the surrounding landscape, and landscape-scale
170 effects have also been documented in some fragmentation experiments [21]. Mensurative
171 fragmentation studies now routinely incorporate data on both patch- and landscape-level
172 forest cover patterns, but none of the existing forest fragmentation experiments have
173 explicitly included landscape forest cover in their designs. To date, the only experiments
174 incorporating such a landscape-scale design feature have been conducted at small spatial
175 scales such as the fractal grass and sand mosaic landscapes generated by With et al. [22].

176

177 Another feature of fragmented environments that has not yet been subjected to a direct
178 controlled experimental test is the effect of the surrounding habitat matrix.
179 Fragmentation reviews have stressed for two decades that the matrix may be the strongest
180 determinant of within-fragment conditions [23-25], and these conclusions have recently
181 been supported by a meta-analysis of the responses of more than 1 000 species to habitat
182 area and isolation [26]. However, actual experimental manipulation of the matrix habitat
183 poses a formidable challenge for a forest fragmentation experiment. All three large-scale
184 fragmentation experiments created fragments that are embedded within a single land use
185 type. Of these, only the BDFFP has been able to test for matrix effects as a *post-hoc* by-
186 product of land abandonment by the cattle ranchers who manage the matrix surrounding

the experimental fragments, rather than as an *a priori* component of the experimental design [17, 27]. At the BDFFP, the initial conversion of forest to cattle pasture that created the fragments was partly supported by government subsidies, but the ranches that were established proved to be unprofitable and were subsequently abandoned. Consequently, the matrix at the BDFFP can now be divided into three categories: cattle pasture, *Vismia* regrowth on abandoned pastures that were burned, and *Cecropia* regrowth on abandoned pastures that were not burned [28].

3. OPPORTUNITIES AND CONSTRAINTS FOR THE SAFE PROJECT

The influence of landscape-level forest cover on the ecological patterns and processes within individual fragments has not yet been subjected to a large-scale experimental test, but will play a central part in the Stability of Altered Forest Ecosystems (SAFE) Project. SAFE will be based in Malaysian Borneo, where the Royal Society South East Asian Rainforest Research Programme (SEARRP, www.searrp.org) has a long-standing relationship with the Sabah Foundation (Yayasan Sabah, www.ysnet.org.my), a state government body charged with spearheading the socio-economic development of the Malaysian state of Sabah. As part of this role, the Sabah Foundation is converting a portion of their forestry estate to oil palm plantation, creating an opportunity for a large-scale deforestation and forest fragmentation experiment. The forest that will be converted has, like most of the forest in the region, already undergone two rounds of selective logging (Table 1). The Sabah Foundation, in collaboration with its subsidiary company Benta Wawasan and with the Sabah Forestry Department, set the time frame of the plantation development such that forest conversion will occur in 2011, giving an

210 opportunity to collect pre-fragmentation data. The long collaborative history between the
211 Sabah Foundation and SEARRP, combined with the high profitability of palm oil [29],
212 work together to ensure the long-term persistence of the experimental fragments that the
213 SAFE Project will create.

214
215 In total, the entire SAFE Project experimental site has an area of 7200 ha. The
216 experimental site currently connects a Virgin Jungle Reserve (VJR) of 2200 ha to a large
217 area of forest (> 1 million ha). Most of the large expanse has been through either one or
218 two rotations of selective logging, although it also encompasses three large conservation
219 areas that have never been logged (Danum Valley, Maliau Basin and Imbak Canyon).
220 The VJR will become isolated during the conversion process. Within the experimental
221 block, the Sabah Foundation agreed to allow up to 800 ha of cultivatable land to be set
222 aside as forest fragments. In addition to this allowance, Malaysian law prohibits the
223 clearance of forest on steep slopes and along permanent streams, accounting for another
224 approximately 500 ha (~7 % of the experimental area). The size of the experimental area,
225 combined with the land that is legally prohibited from clearance and the amount of land
226 that can be retained for experimental purposes makes it feasible to place fragments in
227 locations that will vary in the amount of forest cover in the landscapes surrounding them.
228 This provides a unique opportunity to assess in a controlled, experimental design the
229 impact of landscape context on a wide range of ecological dynamics in forest fragments.

230
231 We stress that the SAFE Project is not the cause of deforestation in this landscape.
232 Rather, it will be embedded within an independent establishment of a large oil palm

plantation in a lowland dipterocarp rainforest. The project is located in an area that has been gazetted for conversion to plantation for the last 20 years, and will use a planned and government approved oil palm conversion to conduct a large-scale landscape experiment. As such, the SAFE Project is using an opportunity that has arisen from the expansion of the oil palm industry to establish an experiment that can simultaneously address basic scientific questions, the answers to which should be important for forest management and conservation in tropical Asia.

3.1 Requirements for the SAFE experimental design

Large-scale experiments must meet the needs of multiple users, both now and in the future. Here, we outline a set of four criteria that the experimental design needs to conform to in order to meet this requirement.

1. For any experiment, it is important to have a relatively standard design that generates well-replicated data in a transparent and open manner. This enables researchers to employ well-established statistical methods that produce unambiguous results and inferences while still allowing supplementary analyses.
2. The BDFFP experience shows that, over the lifetime of a long-term experiment, research teams investigate a wide range of ecological patterns and processes that vary in the spatial scales over which they operate. It is difficult to predict the range of questions and taxa that will ultimately be studied over the long haul of a landscape experiment. But regardless of the details, to obtain maximum benefit from having multiple teams investigating multiple phenomena at a given site, the layout of the experiment and the placement of sampling points need to be designed so as to allow

the investigation of ecological phenomena across varying spatial scales. The design should allow for data on those various phenomena to be tied together, allowing linkages among them to be subjected to statistical analyses. This mandates a form of hierarchically structured, multilevel spatial design for the fragments and the sampling points within them.

3. As an experiment, it will be possible to collect pre-fragmentation data, which greatly increases the power of any analysis. However, as happened at the BDFFP, it is likely that many researchers will only begin to use the SAFE fragments once they have been created. Because these researchers will have no opportunity to collect pre-fragmentation data, it is vital to ensure that the spatial layout of the experiment also permits the use of appropriate spatial controls, as are commonly used in observational studies. Control sites should be located as close as possible to the experimental fragments, yet be located as far as logistically feasible from forest edges. The location of forest control sites does, however, need to be traded off against the logistical difficulties of accessing those sites.

4. Environmental variation can become a major source of experimental error [16]. Consequently, the design should control for obvious and known gradients in environmental variation, such as altitude and slope. Aspect is a less important issue for tropical studies than it is for temperate studies, where there can be substantial differences in biological patterns on north vs south facing slopes [30-31].

4. A FRACTAL SAMPLING DESIGN FOR UNIFYING DATA ACROSS SPATIAL SCALES

We suggest that it is possible to meet the first two criteria presented above by employing a hierarchical sampling design based around a triangular fractal pattern (Fig. 1), and extending that design to define the placement of fragments. To our knowledge, fractal sampling designs have not been used in prior experimental landscape studies, and in general are still scant in empirical ecological investigations.

Triangles were chosen as three sampling points is the minimum required to generate multiple estimates of point-to-point turnover in community composition (β diversity), allowing researchers to estimate the mean and variance of β diversity at a given distance. The distance between sample points at the finest scale of the fractal is determined by a desire to ensure the results from this project are comparable to those of the BDFFP. The distances from the edge to interior of circular fragments of size 1, 10 and 100 ha are respectively $10^{1.75}$, $10^{2.25}$ and $10^{2.75}$ m. Consequently, the 1st-order base of the fractal consists of equilateral triangles with sides of 56 m ($10^{1.75}$) and a sampling point at each vertex. These 1st-order fractals will be placed so that the centroids of the triangles are located on the vertices of a 2nd-order fractal consisting of an equilateral triangle with 178 m sides ($10^{2.25}$). The 2nd-order fractals will be embedded within 3rd- and 4th-order fractals with sides of 564 and 1780 m ($10^{2.75}$ and $10^{3.25}$) respectively (Fig. 1). Consequently, one 4th-order fractal comprises 81 sampling points which are separated by distances distributed evenly along a \log_{10} -scale, providing a total of 81, 243, 729 and 2187 pairwise combinations of traps separated by mean distances of $10^{1.75}$, $10^{2.25}$, $10^{2.75}$ and $10^{3.25}$ m respectively (Fig. 1a).

This sampling design provides a clearly defined structure for aggregating data on ecological phenomena that vary over different spatial scales, paving the way for the unification of data collected on different components of the ecosystem. Ecological patterns that are expected to vary over fine spatial scales, such as microclimate or soil microbial composition, could be sampled at the vertices of the 1st-order fractals. Tree and large mammal communities, by contrast, should vary over larger spatial scales and could be sampled at the vertices of the 2nd or higher-order fractals. At each of the 2nd-order vertices, data collected at the three vertices of the 1st-order fractal could either be aggregated to scale up the data to the same spatial scale as the variables measured at the 2nd-order fractal, or treated as pseudo-replicates of 1st-order phenomena.

5. THE SAFE PROJECT EXPERIMENTAL DESIGN

5.1 Experimental forest fragmentation

The fractal sampling pattern will be used to define the physical location of fragments, with the series of distances used to separate sampling points in the control sites replicated within and among the experimental fragments. The SAFE Project will be a split-plot experiment comprising six replicate blocks (A-F in Fig. 2), each containing four plots with samples in either 4 × 1-ha fragments, 2 × 10-ha fragments, 1 × 100-ha fragment, or in pre-fragmentation continuous forest that will be converted to oil palm (i.e. matrix samples). Each of the six blocks thus contains seven fragments in three size treatments, giving a total of 42 experimental fragments (total area = 744 ha). To be directly comparable with the BDFFP, SAFE will create fragments of 1, 10 and 100 ha. This range of fragment sizes is also very relevant to policy decisions about land-use change, as

the real-world distribution of fragment sizes typically results in more than 90 % of fragments being 100 ha or smaller [32-33]. However, fragments in the SAFE Project will be circular, thereby minimising the edge:area ratio of fragments. This differs from the square plots used at previous forest fragmentation experiments, and ensures the negative impacts of edge effects will be kept to a minimum in this experiment.

Fragments within plots are located to ensure that there is (1) equal sampling effort across plot size classes and (2) an equal spatial distribution of sampling effort in the four plots (Fig. 1), analogous to the design of the Kansas Fragmentation Study [34]. Plots with fragments will be separated by 178 m ($10^{2.25}$ m). The fourth plot with the matrix samples will be aligned with the 100-ha fragment to create a large, 1128 m edge gradient extending from the centre of the fragment, across the fragment edge and out into the matrix. The spatial distribution of sampling points within and among fragments is based on a transect derived from the fractal sampling scheme used in the continuous habitat controls (Fig. 1), ensuring that the distances among samples in the fragments and controls are directly comparable.

The six experimental blocks are placed within the experimental site in a non-random manner in such a way as to maximise the range of forest cover that will remain in the landscapes surrounding sampling points. Average forest cover around sampling points within the six blocks ranges from 16 to 50 % when a landscape is described as a circle with a 3 km radius (Table 2). Experimental blocks are also oriented within the study area in a non-random manner to take advantage of the local topography, such that sampling

transects run at roughly equal altitude. Finally, blocks are also oriented to remove or minimise other potentially confounding effects such as slope, latitude, longitude, and distance to forest edges prior to the forest conversion (see analyses below).

In addition to the sampling sites in the control habitats and experimental fragments, SAFE will take advantage of a 2200 ha Virgin Jungle Reserve that will become isolated during the forest conversion process (Fig. 2). A single transect of sampling points will be placed from the edge to interior of the VJR. The transect extends more than twice the distance of the transects in the experimental fragments, representing the larger size of the VJR, but the distribution of sampling points is still based on the fractal pattern used elsewhere. A similar, long transect will be placed from the edge to interior of the continuous logged forest habitat.

5.2 Control sites and the forest modification gradient

One 4th-order fractal sampling network will be established in each of three control habitats; continuous old growth forest, continuous logged forest and continuous oil palm plantation. Taken together, the control habitats and experimental fragments represent a comprehensive gradient of habitat modification comprising (1) old growth rainforest, (2) logged rainforest, (3) logged and fragmented rainforest, and (4) intensive agriculture in the form of an oil palm plantation. Moreover, each of these levels of habitat modification incorporate different land use and disturbance intensities (Table 1). For example, in the old growth forest sites, located in the Maliau Basin Conservation Area, one of the 3rd order groups of sampling points is located in the water catchment of the Maliau Basin

Field Centre (OG3 on Fig. 1). This water catchment was lightly logged in the 1970s and again in the mid-1990s to provide timber for the field centre, but the vertical structure and species composition of the canopy and undergrowth communities in this area remain representative of primary forest in the wider region. Similarly, the six blocks of experimental fragments vary in the level of logging damage and in the amount of forest cover surrounding them, and the oil palm control sites vary in the age and canopy cover of the palms. The forest modification gradient that is built into the SAFE experimental design mimics the real-world pattern of habitat conversion in Borneo, ensuring that phenomena observed in the study should be directly pertinent to policy issues in the region.

5.3 Experimental variation of riparian corridors

Many tropical nations now have environmental legislation prohibiting the clearance of forest along the banks of streams and rivers. In Malaysia, the legal requirement is for a 30 m strip of forest to be left standing on either side of streams with permanent above-ground water flows. Legislation of this type is designed primarily to protect water quality, and has the added benefit of preserving wildlife corridors connecting standing forest remnants. SAFE will experimentally manipulate the width of riparian corridors to examine their efficacy and with a view to identifying the optimum width for future land use conversions (Fig. 2).

We identified six micro-watersheds within the experimental area that have approximately equal area ($260 \text{ ha} \pm \text{SD } 10$) and slope ($16^\circ \pm \text{SD } 2$). Watersheds of this size contain

headwater streams that are approximately 2 km long. The six watersheds vary in the amount of forest cover that will remain following conversion to oil palm (range 3-30 %; forest cover estimates include the experimental fragments). Each experimental watershed has been assigned one of six riparian widths (0, 5, 15, 30, 60, and 120 m on each side of the stream), with the widths assigned in a way that ensures log-transformed width is not confounded with watershed size ($r_4 = 0.28$, $P = 0.60$), average watershed slope ($r_4 = -0.05$, $P = 0.93$) or forest cover ($r_4 = 0.64$, $P = 0.17$). The permanent streams in the experimental watersheds are small (2 – 3 m across), and are typically enclosed by the forest canopy which meets over the stream. Consequently, from a biodiversity corridors perspective, the range of corridor widths is two times that of the riparian widths (0 – 240 m).

In addition to the experimental watersheds, we have identified three control watersheds (1 × old growth forest, 1 × logged forest and 1 × oil palm plantation). Control watersheds were chosen to match the experimental watersheds as closely as possible in terms of size and slope.

5.4 Analyses of potential confounding factors within the SAFE experimental design

We conducted a series of analyses to ensure that the spatial design of the SAFE experiment will not be confounded with known physical features of the environment. All validation analyses were based on the 2nd-order fractal, meaning that the groups of three sampling points separated by 56 m were represented by a single datapoint. This was because the GIS data used in these analyses are on a 60 m grid, so 1st-order sampling

points often fell within the same grid-square and do not represent separate points for analysis. We modelled the effect of fragment size, transect order (distance along the transect within a plot; Fig. 1), and their interaction on a set of variables that could potentially confound future analyses: (1) latitude and longitude; (2) altitude; (3), slope; (4) pre-fragmentation land use context, represented by distance to forest edges prior to the forest conversion; and (5) isolation, represented by distance to large forest areas following forest conversion (we defined large forest areas as the VJR and the continuous area of logged forest to the north of the experimental area: the VJR is included because much of it has not been logged, and because it is a large area of forest in which temporal changes and species losses are expected to be very slow). Fragment size and transect order were both modelled as categorical variables, with transect order taking the values 1–4 for sample points extending from the centre to the extremity of each plot (Fig. 1). Matrix samples were included in the analysis, and coded as having a fragment ‘area’ of 0 ha. The split-plot experimental design imposes a hierarchical sampling structure on the data, so effects were tested using mixed effects models [35-36] with *p*-values estimated on the basis of Markov Chain Monte Carlo samples (function *aovlmer.fnc* in the R package ‘languageR’ [37]). All analyses were conducted using the ‘lme4’ [38] and ‘languageR’ [37] packages for R.2.12.1 software [39], and the results are summarised in Table 3.

When the location of blocks is accounted for, our analyses show that there should be no significant confounding of latitude with fragment area, transect order or the area × transect interaction. Moreover, the main effects of fragment area, transect order or the

area \times transect interaction will not be confounded with altitude, slope, distance to pre-fragmentation land use or isolation (Table 3).

Mean altitude at sampling points across the experimental blocks is 450 m (median = 460 m, interquartile range 72 m), and mean altitude within blocks ranges from a minimum of 400 m (Block B) to a maximum of 470 m (Block F). Altitude is partially confounded with landscape forest cover: when forest cover is estimated at the 3-, 4- and 5-km scales, there is a strong, negative correlation with altitude ($r < -0.80$, $P < 0.001$). These correlations are driven by the lower altitude of Block B and are not significant if that block is removed from the analysis. Altitude and forest cover at smaller spatial scales are also confounded, although the correlation is not significant ($|r| < 0.53$, $P > 0.27$).

The location of sampling points with respect to pre-fragmentation land use is of concern, as edge effects that are already impacting the experimental area may alter ecological patterns detected in the pre-fragmentation data, reducing the likely impact that experimental habitat fragmentation would have had, compared to a uniform landscape. Experimental blocks will vary in their distance from the pre-fragmentation forest edges, with mean distances ranging from 910 m (Block C) to 2680 m (Block B). All sampling points will be at least 585 m from the pre-fragmentation edges and 87 % of sampling points will be more than 1 km away, which is beyond the expected penetration distance of most edge effects, as found in the BDFFP [8]. As with distance to pre-fragmentation forest edges, the experimental blocks will vary in their isolation distance to large forest areas, with mean isolation distances ranging from 870 m (Block D) to 3130 m (Block C).

463

464 ***5.5 Forest cover in the landscape surrounding experimental blocks***

465 We considered the proportion of the landscape that will remain forested around an
466 individual sampling point following the experimental fragmentation to be landscape
467 forest cover. Blocks will vary significantly in the average amount of landscape forest
468 cover at the four largest spatial scales tested (Table 2a). Blocks D and F are located such
469 that forest cover shows little variation at any spatial scale (33-36 % and 31-34 %
470 respectively), whereas forest cover at Block B increases from 36 % at 1 km scale to
471 almost 60 % at 5 km scale. Across all experimental blocks, forest cover at any spatial
472 scale is tightly correlated with forest cover at similar scales, but that correlation weakens
473 as the difference in spatial scales increases (Table 2c).

474

475 Proportion forest cover in the landscape will be strongly confounded with distance of
476 isolation from large forest areas, with significant correlations existing between isolation
477 distance and forest cover at all spatial scales, and particularly strongly for the 2- and 3-
478 km landscapes ($r < -0.75$, $P < 0.001$ for both correlations). Such correlations are almost
479 inescapable in landscape experiments, and suggest that isolation from large forest areas
480 and landscape forest cover are different metrics reflecting the same landscape-scale
481 gradient in forest cover [19].

482

483 **6. DISCUSSION**

484 Opportunities to design and implement large-scale, long-term ecosystem experiments are
485 rare, but may provide an important method to assess the impacts of global change on

ecosystems [12]. The present-day research and policy environment requires a detailed understanding of the effects of habitat modification on the ability of tropical forests to deliver ecosystem services, such as carbon sequestration and the maintenance of water supplies, and to support the high levels of biodiversity for which they are renowned [7]. Consequently, there is a definable need to establish a new, large-scale forest fragmentation experiment which can address a new generation of research and policy questions. Locating such an experiment in a region undergoing heavy logging and rapid conversion to high-intensity agricultural systems, and designing the experiment to mimic the sequence of land use change and the types of habitats that are being generated by economic forces in the region, ensures that such a project will have direct relevance to high-profile policy issues. Here, we have presented the design for the Stability of Altered Forest Ecosystems Project which is being established in the lowland dipterocarp forests of Malaysian Borneo. Most such experiments have been carried out in temperate or boreal regions [10], and no such large landscape experiment has yet been conducted in tropical Asia. Yet this region is of critical importance, as it harbors a large fraction of the Earth's endangered biodiversity [40-41] and is where the conservation status of threatened species has deteriorated most rapidly in the last three decades [42].

In a southeast Asian context, ongoing deforestation threatens impending biodiversity losses in the region [43], a scenario that has already played out in Singapore [44]. Deforestation has been driven forward as a result of illegal logging activities that have been facilitated by corruption [45], and the logged forests become more susceptible to wild fires [46], which greatly amplify the negative effects of forest loss upon

biodiversity. Moreover, the proliferation of oil palm plantations in Southeast Asia is placing tremendous pressure on forest cover [47-48]. Oil palm is the world's primary source of vegetable oil and fat [49] and one of the world's most rapidly expanding crops [48]. Developing biofuel markets are likely to further increase global demand for palm oil, which generates high yields at low costs [48] creating large profit margins [29]. Oil palm revenue already accounts for almost 2 % of the gross national income of Indonesia and more than 5 % for Malaysia [50]. These economic drivers puts tremendous pressure on the remaining, and shrinking, forests. Part of the stimulus for biofuels is a desire to reduce carbon emissions from fossil fuels, but total carbon emission reductions are unlikely to be achieved when tropical forests are cleared to make way for oil palm plantations [48, 51].

The SAFE Project will be embedded in a planned conversion of logged, lowland dipterocarp forest to oil palm plantation. Over the duration of the experiment, the matrix surrounding the forest fragments will change as the plantation progresses from a cleared forest through the process of terracing, planting of cover crops and oil palm, and the gradual maturation of oil palm which should form a canopy approximately seven years after planting. Although the matrix will change through time, at any given point in time all the fragments will be surrounded by a matrix in a similar state. This means that it will not be possible to experimentally test matrix effects in this project. Nonetheless, SAFE represents substantial advances over existing forest fragmentation experiments. Perhaps most importantly, SAFE represents the first opportunity to experimentally test the effects of landscape forest cover on ecological patterns within isolated patches. By locating the

experimental blocks in positions to maximise the range of naturally-occurring forest cover in the wider landscape surrounding the experimental fragments, we allow for direct tests of the ways in which landscape forest cover may moderate patch-level effects. This has been achieved without sacrificing the ability to detect patch-level effects, and while ensuring that results from this project will be directly comparable to those of existing experimental forest fragmentation studies. Moreover, the unification of a hierarchical, fractal-based sampling scheme with the spatial layout of the fragments will allow greater integration of data collected on ecological patterns and processes that operate over very different spatial scales. The riparian component of the project will help resolve the question of how to design effective corridors so as to protect water resources and aquatic biodiversity. Finally, the use of three control habitats inserts the experimentally fragmented landscape into a wider gradient of habitat modification and land use intensity, allowing for a much broader understanding of the biotic and abiotic impacts of land use change.

The reality of a ‘natural ecosystem’ is fast disappearing as humans modify the world at an ever-accelerating pace, meaning much of the world’s biodiversity must now perish or persist in human-modified landscapes [7]. Understanding how much of the diversity of life will persist in the future will require a better understanding of ecological processes that occur when remnants of natural habitats are embedded in a matrix dominated by human activities [52]. The SAFE Project will be one of the world’s largest ecological experiments and is designed to directly address questions about how logging, forest fragmentation and deforestation modify the functioning of tropical rainforests, impair

their ability to deliver ecosystem services, and reduce their capacity to support the diversity of life.

Acknowledgements

We thank the Sime Darby Foundation, the Sabah Foundation, Benta Wawasan, and the Sabah Forestry Department for their generous support of the SAFE Project. Mick Crawley, Charles Godfray, Bill Laurance, Georgina Mace and Stuart Pimm provided valuable advice on the project design. This paper arose from a workshop sponsored by the NERC Centre for Population Biology. Researchers are invited to take advantage of the SAFE study sites and, if interested, should visit the project website (www.safeproject.net).

References

1. Meffe GK, Carroll CR. Principles of conservation biology. 2nd ed. Sunderland: Sinauer Associates; 1997.
2. MacArthur RH, Wilson EO. The theory of island biogeography. Princeton: Princeton University Press; 1967.
3. Simberloff D, Abele LG. Island biogeography theory and conservation practice. Science. 1976;191:285-6
4. Diamond JM. The island dilemma: Lessons of modern biogeographic studies for the design of natural reserves. Biol Conserv. 1975;7:129-46
5. Lovejoy TE, Bierregaard RO, Rankin JM, Schubart HOR. Ecological dynamics of tropical forest fragments. In: Sutton SL, Whitmore TC, Chadwick AC, eds. *Tropical rain*

578 *forest: Ecology and management*. Oxford: Blackwell Scientific Publications; 1983:377-
579 84.

580 6. Lovejoy TE, Oren DC. The minimum critical size of ecosystems. In: Burgess RL,
581 Sharpe DM, eds. *Forest island dynamics in man-dominated landscapes*. New York,
582 USA: Springer-Verlag; 1981:7-12.

583 7. Gardner TA, Barlow J, Chazdon RL, Ewers RM, Harvey CA, Peres CA, et al.
584 Prospects for tropical forest biodiversity in a human-modified world. *Ecol Lett*.
585 2009;12:561-82

586 8. Laurance WF, Lovejoy TE, Vasconcelos HL, Bruna EM, Didham RK, Stouffer PC, et
587 al. Ecosystem decay of amazonian forest fragments: A 22-year investigation. *Conserv*
588 *Biol*. 2002;16(3):605-18

589 9. Block WM, Franklin AB, Ward JP, Ganey JL, White GC. Design and implementation
590 of monitoring studies to evaluate the success of ecological restoration on wildlife.
591 2001;9(3):293-303

592 10. Debinski DM, Holt RD. A survey and overview of habitat fragmentation experiments.
593 *Conserv Biol*. 2000;14(2):342-55

594 11. Schmiegelow FKA, Machtans CS, Hannon SJ. Are boreal birds resilient to forest
595 fragmentation? An experimental study of short-term community responses. *Ecology*.
596 1997;78(6):1914-32

597 12. Clark JS, Carpenter SR, Barber M, Collins S, Dobson A, Foley JA, et al. Ecological
598 forecasts: An emerging imperative. *Science*. 2001 July 27, 2001;293(5530):657-
599 60.(10.1126/science.293.5530.657)

600 13. Gilbert-Norton L, Wilson R, Stevens JR, Beard KH. A meta-analytic review of
601 corridor effectiveness. *Conserv Biol.* 2010 Jun;24(3):660-68.(10.1111/j.1523-
602 1739.2010.01450.x)

603 14. Haddad NM. Corridor and distance effects on interpatch movements: A landscape
604 experiment with butterflies. 1999;9(2):612-22

605 15. Damschen EI, Haddad NM, Orrock JL, Tewksbury JJ, Levey DJ. Corridors increase
606 plant species richness at large scales. 2006 September 1, 2006;313(5791):1284-
607 86.(10.1126/science.1130098)

608 16. Margules CR. The wog wog habitat fragmentation experiment. *Environ Conserv.*
609 1992;19(4):316-25

610 17. Stouffer PC, Strong C, Naka LN. Twenty years of understorey bird extinctions from
611 amazonian rain forest fragments: Consistent trends and landscape-mediated dynamics.
612 *Divers Distrib.* 2009;15(1):88-97

613 18. Joppa LN, Loarie SR, Pimm SL. On the protection of "Protected areas". *PNAS.* 2008
614 May 6, 2008;105(18):6673-78.(10.1073/pnas.0802471105)

615 19. Fahrig L. Effects of habitat fragmentation on biodiversity. *Annu Rev Ecol Evol Syst.*
616 2003;34:487-515

617 20. Donovan TM, Jones PW, Annand EM, Thompson FR, III. Variation in local-scale
618 edge effects: Mechanisms and landscape context. *Ecology.* 1997;78(7):2064-75

619 21. Schweiger EW, Diffendorfer JE, Pierotti R, Holt RD. The relative importance of
620 small-scale and landscape-level heterogeneity in structuring small mammal distributions:
621 An experimental study of habitat fragmentation. In: Barrett G, Peles J, eds. *Landscape*
622 *ecology of small mammals.* New York: Springer-Verlag; 1999.

- 623 22. With KA, Cadaret SJ, Davis C. Movement responses to patch structure in
624 experimental fractal landscapes. *Ecology*. 1999;80(4):1340-53
- 625 23. Saunders DA, Hobbs RJ, Margules CR. Biological consequences of ecosystem
626 fragmentation: A review. *Conserv Biol*. 1991;5(1):18-32
- 627 24. Ewers RM, Didham RK. Confounding factors in the detection of species responses to
628 habitat fragmentation. *Biol Rev*. 2006;81(1):117-42
- 629 25. Kupfer JA, Malanson GP, Franklin SB. Not seeing the ocean for the islands: The
630 mediating influence of matrix-based processes on forest fragmentation effects. *Global*
631 *Ecol Biogeog*. 2006;15:8-20
- 632 26. Prugh LR, Hodges KE, Sinclair ARE, Brashares JS. Effect of habitat area and
633 isolation on fragmented animal populations. *PNAS*. 2008 December 30,
634 2008;105(52):20770-75.(10.1073/pnas.0806080105)
- 635 27. Nascimento HEM, Andrade ACS, Camargo JLC, Laurance WF, Laurance SG, Ribeiro
636 JE. Effects of the surrounding matrix on tree recruitment in amazonian forest fragments.
637 *Conserv Biol*. 2006;20(3):853-60
- 638 28. Mesquita RCG, Ickes K, Ganade G, Williamson GB. Alternative successional
639 pathways in the amazon basin. 2001;89(4):528-37.(10.1046/j.1365-2745.2001.00583.x)
- 640 29. Fisher B, Edwards DP, Giam X, Wilcove DS. The high costs of conserving southeast
641 asia's lowland rainforests. in press:null.(doi:10.1890/100079)
- 642 30. Brothers TS, Spingarn A. Forest fragmentation and alien plant invasion of central
643 indiana old-growth forests. *Conserv Biol*. 1992;6(1):91-100
- 644 31. McDonald RI, Urban DL. Forest edges and tree growth rates in the north carolina
645 piedmont. *Ecology*. 2004;85(8):2258-66

646 32. Cochrane MA, Laurance WF. Fire as a large-scale edge effect in amazonian forests. J
647 Trop Ecol. 2002;18:311-25

648 33. Ribeiro MC, Metzger JP, Martensen AC, Ponzoni FJ, Hirota MM. The brazilian
649 atlantic forest: How much is left, and how is the remaining forest distributed?
650 Implications for conservation. Biol Conserv. 2009 Jun;142(6):1141-
651 53.(10.1016/j.biocon.2009.02.021)

652 34. Holt RD, Robinson GR, Gaines MS. Vegetation dynamics in an experimentally
653 fragmented landscape. Ecology. 1995;76(5):1610-24

654 35. Sokal RR, Rohlf FJ. Biometry. 3rd Edition ed. New York: W. H. Freeman and
655 Company; 1995.

656 36. Pinheiro JC, Bates DM. Mixed-effects models in s and s-plus. New York: Springer;
657 2000.

658 37. Baayen RH. Languager: Datasets and functions with "Analyzing linguistic data: A
659 practical introduction to statistics". R package version 1.0; 2010.

660 38. Bates D, Bolker B. Lme4: Linear mixed-effects models using s4 classes. R package
661 version 0.999375-39; 2011.

662 39. R Development Core Team. R: A language and environment for statistical computing.
663 Vienna: R Foundation for Statistical Computing; 2011.

664 40. Myers N, Mittermeier CG, Mittermeier RA, Fonseca GAB, Kent J. Biodiversity
665 hotspots for conservation priorities. Nature. 2000;403:853-58

666 41. Orme CDL, Davies RG, Burgess M, Eigenbrod F, Pickup N, Olson VA, et al. Global
667 hotspots of species richness are not congruent with endemism or threat. 2005;436:1016-
668 19

669 42. Hoffmann M, Hilton-Taylor C, Angulo A, Böhm M, Brooks TM, Butchart SHM, et
670 al. The impact of conservation on the status of the world's vertebrates. 2010 December
671 10, 2010;330(6010):1503-09.(10.1126/science.1194442)

672 43. Sodhi NS, Pin Koh L, Brook BW, Ng PKL. Southeast asian biodiversity: An
673 impending disaster. Trends Ecol Evol. 2004;19(12):654-60

674 44. Brook BW, Sodhi NS, Ng PKL. Catastrophic extinctions follow deforestation in
675 singapore. Nature. 2003;424:420-23

676 45. Smith J, Obidzinski K, Subarudi, Suramenggala I. Illegal logging, collusive corruption
677 and fragmented governments in kalimantan, indonesia. Int For Rev. 2003;5(3):293-302

678 46. Langner A, Siegert F. Spatiotemporal fire occurrence in borneo over a period of 10
679 years. Global Change Biol. 2009;15(1):48-62

680 47. Koh LP, Wilcove DS. Is oil palm agriculture really destroying tropical biodiversity?
681 Conserv Lett. 2007;1:60-64

682 48. Fitzherbert EB, Struebig MJ, Morel A, Danielsen F, Brühl CA, Donald PF, et al. How
683 will oil palm expansion affect biodiversity? Trends Ecol Evol. 2008;23(10):538-45

684 49. Turner EC, Snaddon JL, Fayle TM, Foster WA. Oil palm research in context:
685 Identifying the need for biodiversity assessment. PLoS ONE. 2008;3(2):e1572

686 50. Koh LP, Wilcove DS. Cashing in palm oil for conservation. Nature.
687 2007;448(7157):993-94

688 51. Fargione J, Hill J, Tilman D, Polasky S, Hawthorne P. Land clearing and the biofuel
689 carbon debt. Science. 2008 February 29, 2008;319(5867):1235-
690 38.(10.1126/science.1152747)

691 52. Rosenzweig MR. Win-win ecology: How the earth's species can survive in the midst
692 of the human enterprise. New York: Oxford University Press; 2003.
693
694

695 Table 1: The land use intensity gradient incorporated into the SAFE Project, extending
 696 from primary forest through to oil palm plantation. Blocks correspond to the sampling
 697 sites illustrated in Fig. 2 and are ordered from least to most disturbed.
 698

Block	Habitat	Logging	Fragmentation*	Forest cover [†]	Forest quality (range) [‡]	Notes
OG1	Forest	Never	Continuous	100 %	4.44 (3-5)	> 1 km from reserve boundary
OG2	Forest	Never	Continuous	100 %	4.88 (4-5)	> 500 m from reserve boundary
OG3	Forest	Low intensity	Continuous	100 %	4.22 (3-5)	Lightly logged 1970s and 1990s
LF1	Forest	Twice	Continuous	100 %	3.22 (3-4)	Twice logged
LF3	Forest	Twice	Continuous	76 %	3.44 (3-4)	Twice logged
LF2	Forest	Twice	Continuous	86 %	3.67 (3-4)	Twice logged
LFE	Forest	Twice	Continuous	71 %	3.25 (2-4)	Twice logged
VJR	Forest	Variable	Fragmented	61 %	3.43 (2-5)	Logged around edges, never logged in the steep interior
B	Forest	Twice	Fragmented	50 %	2.75 (2-4)	Twice logged
D	Forest	Twice	Fragmented	35 %	2.06 (1-3)	Twice logged
F	Forest	Twice	Fragmented	34 %	2.50 (1-3)	Twice logged
A	Forest	Twice	Fragmented	26 %	2.25 (1-4)	Twice logged
E	Forest	Twice	Fragmented	21 %	1.94 (1-4)	Twice logged
C	Forest	Twice	Fragmented	16 %	2.06 (1-4)	Twice logged
OP3	Oil palm	NA	Cleared	15 %	NA	Planted 2000; closed canopy; some cover crop; 1 km from forest
OP2	Oil palm	NA	Cleared	40 %	NA	Planted 2006; canopy just forming; cover crop; 500 m from forest
OP1	Oil palm	NA	Cleared	15 %	NA	Planted 2006; canopy just forming; cover crop; 700 m from forest

699 *Fragmentation: Continuous forest cover means the sampling block is located in a contiguous
 700 forest management area of approximately one million ha.

701 [†] Forest cover: the average amount of forest cover in a 3-km radius surrounding 2nd order
702 sampling points within each block. No distinction is made between primary and logged forest.
703 [‡] Forest quality: average forest quality at 2nd order sampling points within each block. Quality is
704 scored on a qualitative scale of 1-5; (1) very poor, no standing trees, open canopy with ginger,
705 vines or low scrub; (2) poor, open canopy with occasional small trees over a ginger and vine
706 layer; (3) okay, small trees abundant and canopy at least partially closed; (4) good, lots of trees
707 including some large trees and a closed canopy; (5) very good, no evidence of logging, closed
708 canopy with large trees.
709

710 Table 2. Variation in the amount of forest cover surrounding experimental blocks in the
711 SAFE Project. (a) Values represent the *F* and *P*-values from a mixed effect model testing
712 for different forest cover amounts among the experimental blocks at five spatial scales.
713 (b) Mean forest cover (± 1 SE) surrounding sampling points in each of the six
714 experimental blocks. (c) Pearson correlation coefficient between forest cover at the five
715 spatial scales. Values in **bold** are significant with $P < 0.05$.
716

Landscape scale	(a) Block		(b) Forest cover (%)						(c) Correlation			
	$F_{(5,18)}$	<i>P</i>	A	B	C	D	E	F	2 km	3 km	4 km	5 km
1 km	0.78	0.356	34(1)	36(3)	26(2)	36(3)	33(1)	33(2)	0.90	0.79	0.55	0.39
2 km	7.15	0.003	26(1)	36(2)	14(1)	33(1)	21(1)	32(2)		0.92	0.69	0.51
3 km	31.31	<0.001	26(1)	50(1)	16(1)	35(1)	21(0)	34(1)			0.92	0.80
4 km	44.36	<0.001	26(1)	57(1)	26(1)	33(1)	26(0)	32(1)				0.97
5 km	59.23	<0.001	26(1)	59(1)	34(1)	35(0)	30(0)	31(0)				

717

718 Table 3. The potential of six variables to confound future analyses of data emerging from
 719 the SAFE Project. Values represent the *F* and *P*-values from a mixed effect model.

720 Significant relationships are in **bold**.

721

Variable	Block		Fragment area		Transect order		Area × Transect	
	<i>F</i> _(5,18)	<i>P</i>	<i>F</i> _(3,15)	<i>P</i>	<i>F</i> _(3,60)	<i>P</i>	<i>F</i> _(9,60)	<i>P</i>
Longitude	502.279	<0.001	1.102	0.944	0.007	0.998	1.657	0.447
Latitude	75.906	<0.001	0.155	0.998	0.060	0.932	0.205	0.999
Altitude	1.136	0.233	1.432	0.796	0.137	0.514	0.686	0.859
Slope	0.123	0.930	0.820	0.753	0.713	0.570	0.287	0.982
Pre-fragmentation land use	19.377	<0.001	1.708	0.927	0.020	0.345	1.806	0.303
Isolation	24.530	<0.001	0.906	0.991	0.548	0.861	1.340	0.558

722

Figure captions

Figure 1. (a) Fractal geometry of the sampling network in a continuous habitat. Points on the vertices of the white triangles represent sampling locations, and triangles of progressively darker shades indicate a progression from the first to fourth order of the fractal pattern respectively. The box on the lower right encompasses a sampling transect that is used as the basis of the sampling scheme in the forest fragments. (b) Spatial layout of fragments forming a single block in the split-plot experimental design of the SAFE Project, showing how the fractal sampling scheme is embedded within each fragment (circles) and the surrounding matrix. The experimental block is formed of four plots: (1) 1×100 -ha fragment; (2) 2×10 -ha fragments; (3) 4×1 -ha fragments; and (4) samples in the matrix adjacent to the 100-ha fragment. Total sampling effort, and the spatial distribution of sampling effort, is exactly the same within all plots. Numbers 1-4 represent the 'transect order' of sampling points within plot, which is used in analyses of the potential impact of confounding variables.

Figure 2. Map of the Stability of Altered Forest Ecosystems Project, located in Malaysian Borneo. The project encompasses a gradient of forest modification encompassing (a) old growth and lightly logged forest; (b) continuous, twice-logged forest; (c) twice-logged and fragmented forest in an oil palm matrix; and (d) continuous oil palm plantation. (e) The fragmentation experiment comprises six blocks (A-F), each with seven fragments arranged as shown in Fig. 1. In addition to the experimental fragments themselves, sampling across edge gradients will occur in a Virgin Jungle Reserve (VJR) adjacent to the experimental area and at the edge of continuous logged forest (LFE) to the north of

the experimental area. Fragments are currently embedded in a twice-logged forest landscape that will be converted to oil palm as part of the experiment (labeled ‘Oil palm (future)’). The riparian corridor experiment comprises six watersheds located within the experimental area. Sampling points show the locations of the 2nd order fractals across the study area. The OG3 block of control sites in Maliau Basin falls within the water catchment of the Maliau Basin Field Centre and has been selectively logged.

753

FIGURE 1

754




FIGURE 2

